

News 3. Semaglutide 2.4:
novità nel diabete e
nell'obesità

Mikiko Watanabe M.D. Ph.D.



Disclosure

Funding from the following companies:

For providing educational sessions

-Novo Nordisk, Eli Lilly, Theras Lifetech

Institutional research grant support or funding for clinical trials


-Novo Nordisk and Boehringer Ingelheim

Institutional Scientific Board and/or consulting

- Novo Nordisk, Eli Lilly

STEP UP trial design

Randomised, double-blind, placebo- and active-controlled, multinational trial



Population

- Adults (≥18 years)
- BMI ≥30 kg/m²
- ≥1 self-reported unsuccessful dietary effort to lose weight
- Without T2D (HbA_{1c} <6.5%)

Coprimary endpoints*

- Change in bodyweight (%)
- Achieving ≥5% WL

Confirmatory secondary endpoints

- Achieving ≥10%, ≥15%, ≥20%, and ≥25% WL*
- Change in waist circumference (cm)*
- Change in bodyweight (%)[†]
- Achieving ≥20% and ≥25% WL[†]


*Semaglutide s.c. 7.2 mg versus placebo. [†]Semaglutide s.c. 7.2 mg versus 2.4 mg.

BMI=body mass index. HbA_{1c}=glycated haemoglobin. s.c.=subcutaneous. T2D=type 2 diabetes. WL=weight loss. Additional details included in the speaker notes.

Adapted from Supplementary Figure 1: Trial design.

Change in body weight (%)

STEP UP




Data are for the full analysis set, and observed data are from the on-treatment observation period. Treatment comparisons were estimated using the trial product estimand. Error bars are 95% CIs. *Estimated mean change at week 72.
CI, confidence interval; ETD, estimated treatment difference.
Wharton S et al. Lancet Diabetes Endocrinol 2025; DOI: 10.1016/S2213-8587(25)00226-8.

Categorical body weight loss

STEP UP

Trial product estimand




Data are for the full analysis set, and observed data are from the on-treatment observation period. Treatment comparisons were estimated using the trial product estimand. Comparisons of semaglutide 7.2 mg versus 2.4 mg reported in this figure for the proportion of participants with body weight reductions of 5% or greater, 10% or greater, and 15% or greater were conducted in a post-hoc manner. Post hoc analyses were not controlled for multiplicity, and findings for these endpoints should not be used to infer definitive treatment effects.

CI, confidence interval; OR, odds ratio.

Wharton S et al. Lancet Diabetes Endocrinol 2025; DOI: 10.1016/S2213-8587(25)00226-8.

Obesity is a disease that leads to cardio-renal-metabolic complications



OBEesity Treatment Based on Patient Phenotyping

FIRST-LINE THERAPY IS LIFESTYLE MODIFICATION (MEDICAL NUTRITIONAL APPROACH AND IMPLEMENTATION OF PHYSICAL ACTIVITY AND BEHAVIOURAL THERAPY)*

EXCLUDE ENDOCRINE FORMS OF OBEsITY, if suspected, investigate MONOGENIC FORMS

ASSESSMENT OF THE PRESENCE OF COMPLICATIONS AND PATIENT PHENOTYPING

OBEsITY TREATMENT BASED ON PATIENT COMPLICATIONS AND PHENOTYPING


METABOLIC CARDIO RENAL COMPLICATIONS						MECHANICAL COMPLICATIONS		WEIGHT LOSS TARGETS /OWCMC**	AGE	EATING BEHAVIOUR	SPECIAL POPULATIONS	COSTS /QALY
+ASCVD	Indicators of High CV Risk:	HF	CKD	GLYCEMIC CONTROL	MASH/MASLD	OSTEOARTRITIS	OSAS					
Cardiovascular disease was defined as a previous myocardial infarction, previous stroke, or symptomatic peripheral arterial disease in people who are ≥ 45 years of age, with a BMI ≥ 27 kg/m².	Definitions vary; most comprise ≥ 40 years of age, BMI ≥ 27 kg/m², male gender, high blood pressure, atherogenic dyslipidemia, smoking, prediabetes, ASCVD risks (ACC/AHA).	Clinical heart failure (NYHA II–IV) with LVEF ≥50% (tirzepatide) or ≥45% (semaglutide), BMI ≥30 kg/m², reduced functional capacity (KCCQ-CSS <80 or <90), 6-minute walk ≥100 m, and elevated filling pressures (NT-proBNP, echocardiographic criteria, or recent HF hospitalization). T2D was all in the tirzepatide trial.	eGFR < 60 ml/ min per 1.73 m² OR albuminuria (ACR ≥ 3.0 mg/mmol (30mg/g)	PREDIABETES (HbA1c ≥ 5.7% and < 6.5%, FPG > 100 ng/dl < 126 mg/dl, OGTT 120 min glucose ≥ 140 mg/dl < 200 mg/dl) Semaglutide 2.4 mg²² Orlistat 120 mg¹⁴ Metformin: off-label; may reduce diabetes incidence ²⁴ .	Adults with biopsy-confirmed MASH, fibrosis stage (F2–F3) for semaglutide 2.4 and tirzepatide 15 mg, F0-F3 for liraglutide 3 mg , BMI ≥25 kg/m² for semaglutide 2.4 mg and liraglutide 3 mg . BMI ≥27 kg/m² for tirzepatide 15 mg; with or without type 2 diabetes but not for semaglutide 2.4.	Adults ≥18 years with BMI ≥30 kg/m², clinically and radiographically confirmed knee osteoarthritis (ACR criteria; Kellgren–Lawrence grade 2–3), and baseline knee pain ≥40/100 on the WOMAC pain scale.	Adults ≥18 years, BMI ≥30 kg/m², with moderate-to-severe obstructive sleep apnea (AHI ≥15 events/hour); CPAP use allowed (tirzepatide) or excluded (liraglutide); no diagnosis of T2D.	Set individualized weight management goals 5-10% WL Orlistat¹⁴ Liraglutide 3 mg⁷⁴ Naltrexone/ Bupropion²⁷ Phentermine/Topiramate (low-mid doses)⁶⁰	≥ 2 y.o. ¹⁰⁴ Setmelanotide (rare genetic obesity) Metreleptin (rare genetic obesity) 6-12 y.o. Liraglutide 3mg ¹⁰⁹ (EMA only) 12-18 y.o. Semaglutide 2.4mg¹¹³ Phentermine/Topiramate¹¹⁵ (FDA only) Orlistat 120 mg¹¹⁴ (FDA only) 18-65 y.o. Tirzepatide 15 mg Semaglutide 2.4 mg Naltrexone/Bupropion^{27,28} Phentermine/Topiramate²⁶ Orlistat¹⁴ >65 y.o. Semaglutide 2.4 mg ³ (Use with caution; no dose adjustment needed) Tirzepatide 15 mg (Usable; monitor sensitivity in elderly) Liraglutide 3 mg ¹²⁴ (Usable; limited data >75 yrs) Naltrexone/Bupropion ²⁷ (Caution; limited data ≥65 yrs) Phentermine/Topiramate ²⁶ (Caution; monitor CNS/renal) Orlistat ¹⁴ (Usable; limited efficacy data) Sarcopenic obesity Supervised Resistance Training + Protein Intake ≥1.2–1.5 g/kg/day ¹¹⁸ Liraglutide 3 mg is preferred for gradual weight loss with lean mass preservation and proven cardiovascular safety ³³	EMOTIONAL EATING Bupropion / Naltrexone ²⁸ Tirzepatide 15 mg ¹²⁵ BINGE EATING Not AOMs approved explicitly for binge eating disorder. Liraglutide 3.0 mg ¹²³ Phentermine/Topiramate ¹²⁸ Naltrexone/Bupropion ²⁸ Orlistat 120 mg ¹²⁹ Semaglutide 2.4 mg¹²⁵ Tirzepatide 15 mg^{128,134}	CANCER safety No obesity-related risk from Semaglutide 2.4 mg ³ Tirzepatide 15 mg ¹⁴ Evidence from meta-analysis suggests long-term GLP-1 RA use may increase thyroid cancer risk in mixed T2D populations. ¹⁴⁵ No evidence of increased cancer risk (for other AOMs: Orlistat, Phentermine/Topiramate, Bupropion/Naltrexone) Familial Partial Lipodystrophy 1 -3 (FPLD1-3) Acquired Partial Lipodystrophy (Barraquer-Simons) HIV-Associated Lipodystrophy Semaglutide 2.4 mg¹⁵² MONOGENIC OR SYNDROMIC OBEsITY Congenital leptin deficiency Metreleptin¹⁵³ (biallelic POMC, PCSK1, or LEPR deficiency and for BBS) Setmelanotide 3 mg¹⁵⁴ ACQUIRED HYPOTALAMIC OBEsITY Setmelanotide 3 mg¹⁵⁴ Semaglutide 2.4 mg³	BS 176,177
Semaglutide 2.4 mg	Tirzepatide 15 mg^{2,10} Semaglutide 2.4 mg^{3,4} Liraglutide 3 mg¹²	Tirzepatide 15 mg¹¹ Semaglutide 2.4 mg⁵	(use if eGFR>30 ml/min; If eGFR <30 use with caution due to limited data) Orlistat (caution in obesity due to risks of nephrolithiasis or AKI) ¹⁸ Naltrexone/ Bupropion (not recommended) ¹⁵	Semaglutide 2.4 mg²⁹ SGLT2is support weight loss ^{37,38,39,40,41} . If glycemic control allows, insulin ^{44,45} , thiazolidinediones, and sulfonylureas should be avoided ²⁰³ TYPE 1 DIABETES (OFF-LABEL only for weight loss purposes) Semaglutide 2.4 mg⁵¹	Semaglutide 2.4 mg Tirzepatide 15 mg⁵⁶ Liraglutide 3 mg OD⁵⁷	Semaglutide 2.4 mg Tirzepatide 15 mg (NCT01985165)⁶⁸ Orlistat 120 mg¹⁴ Naltrexone/bupropione⁶⁴ Phentermine/Topiramate⁶⁵	Tirzepatide 15 mg⁶⁸ Liraglutide 3 mg⁶⁹ Phentermine/Topiramate⁷⁰	10-20% WL Tirzepatide 5 mg² Semaglutide 2.4mg⁴ Phentermine/Topiramate⁶⁰ >20% WL Tirzepatide 15mg² Semaglutide 7.2 mg⁷⁷ (not available yet) Cagrisema⁷⁸ (not available yet)	Semaglutide 2.4 mg¹²⁵ Tirzepatide 15 mg^{128,134}	Setmelanotide 3 mg¹⁵⁴ Semaglutide 2.4 mg³	PREGNANCY/WILLINGNESS TO CONCEIVE AOMs are either contraindicated or advised against during pregnancy due to insufficient safety data and potential risks to the fetus Stop Liraglutide 3 mg/Semaglutide 2.4 mg at least 2 months before Stop Tirzepatide 15 mg at least 1 month before	

If additional weight loss and reduction of weight-related complications are needed

Bariatric Surgery¹⁷⁷

If insufficient weight loss or weight regain after BS consider: Tirzepatide¹⁹⁷, Semaglutide¹⁹⁴, Liraglutide^{195,196}

*This framework promotes a phenotype-based, individualized approach to obesity care, prioritizing complications and patient preferences over rigid treatment hierarchies, with patient-HCP shared decision-making considering goals, tolerability, contraindications, and cost. **OWCMC, obesity without clinically manifest complications




+ASCVD

Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥ 45 years and with a BMI ≥ 27 kg/m²

Semaglutide 2.4 mg⁴

Naltrexone/
Bupropion
(not recommended)¹⁵

Semaglutide 2.4 mg has shown significant MACE reduction, as well as wider CKM benefits, in people with overweight or obesity and CVD, without diabetes



1. Lincoff AM, et al. N Engl J Med. 2023;DOI:10.1056/NEJMoa2307563.

+ASCVD

Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥45 years and with a BMI ≥27 kg/m²


Semaglutide 2.4 mg ⁴

Naltrexone/
Bupropion
(not recommended)¹⁵

Time from randomisation to first MACE by body weight loss at week 20
In the semaglutide arm

MACE risk reduction with semaglutide was independent of weight loss


Deanfield J et al. *Lancet* 2025;doi:10.1016/S0140-6736(25)01375-3.



Patients at risk (censored)

Placebo									
Weight loss ≥5%	853 (0)	841 (12)	807 (34)	792 (45)	775 (55)	650 (63)	509 (70)	378 (81)	186 (84)
Weight loss ≥0% to <5%	4478 (0)	4407 (51)	4321 (114)	4245 (151)	4166 (195)	3625 (244)	2916 (287)	2067 (324)	932 (341)
Weight gain	3470 (0)	3433 (27)	3363 (75)	3291 (116)	3228 (152)	2868 (193)	2263 (225)	1639 (254)	609 (271)

Time from randomisation to first MACE by body weight loss at week 20
In the placebo arm



Patients at risk (censored)

Semaglutide									
Weight loss ≥5%	5392 (0)	5346 (30)	5265 (79)	5181 (130)	5074 (199)	4454 (245)	3569 (291)	2655 (316)	1197 (339)
Weight loss ≥0% to <5%	2921 (0)	2887 (21)	2825 (65)	2781 (92)	2726 (119)	2413 (142)	1924 (164)	1340 (176)	517 (186)
Weight gain	490 (0)	488 (1)	472 (9)	468 (10)	459 (16)	404 (24)	306 (30)	203 (34)	71 (36)

+ASCVD

HOPE Study Design

Danish hospital records

Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥ 45 years and with a BMI ≥ 27 kg/m²

Exclusion criteria in the 10 years before the index date included diabetes and bariatric surgery

Semaglutide group (semaglutide prescription)

Non-semaglutide group (no semaglutide prescription)

Matched 1:3 on baseline demographics and clinical characteristics

Index date

(date of first filled semaglutide prescription; same date for matched controls)

Follow-up

until end of data availability (8 Dec 2024), migration or outcome of interest

End of study



Inclusion criteria



Aged ≥ 45 years



Hospital diagnosis of overweight or obesity (BMI ≥ 25 kg/m²) in the Danish National Patient Register during index period (1 Jan 2012–31 Dec 2022)



For the semaglutide group, ≥ 1 prescription for once-weekly semaglutide for weight management in Danish Prescription Register (12 Dec 2022–31 Oct 2024)



Outcomes

- Time to 5P-MACE^b
- Time to 3P-MACE^c



Statistics

Kaplan-Meier analysis to generate survival curves and Cox proportional hazards model to generate HRs and 95% CIs

Semaglutide 2.4 mg ⁴

Naltrexone/
Bupropion
(not recommended)¹⁵

^aMatching was based on date of birth, sex, region of residence, educational level (highest attained), obesity category, time between overweight/obesity diagnosis and index date, comorbidities (HF, MI, stroke), predicted CVD risk score and history of ASCVD. ^b5P-MACE was a composite of MI, stroke, hospitalization for HF, coronary revascularization and all-cause mortality. ^c3P-MACE was a composite of MI, stroke and all-cause mortality. 3P-MACE, 3-point major adverse cardiovascular events; 5P-MACE, 5-point major adverse cardiovascular events; ASCVD, atherosclerotic cardiovascular disease; BMI, body mass index; CI, confidence interval; CVD, cardiovascular disease; HF, heart failure; HR, hazard ratio; MI, myocardial infarction.

+ASCVD


Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥ 45 years and with a BMI ≥ 27 kg/m²



Semaglutide 2.4 mg ⁴

Naltrexone/
Bupropion
(not recommended)¹⁵

Semaglutide treatment was associated with relatively lower risk of 3P-MACE^a



Incidence of 3P-MACE per 1000 person-years	
Semaglutide	7.4
Non-semaglutide	14.0
Absolute risk reduction	6.6

Individuals at risk (event rate [%])

Semaglutide group	7113 (–)	6580 (0.20)	5728 (0.31)	4873 (0.39)	4151 (0.53)	3383 (0.67)	2506 (0.77)	1178 (–)
Non-semaglutide group	21,339 (–)	19,704 (0.39)	17,127 (0.65)	14,495 (1.00)	12,333 (1.22)	10,059 (1.38)	7472 (1.48)	3528 (–)

Shaded areas represent the 95% CIs.^a3P-MACE was a composite of MI, stroke and all-cause mortality.
3P-MACE, 3-point major adverse cardiovascular events; CI, confidence interval; HR, hazard ratio; MI, myocardial infarction.

Maximum
follow-up
was 23.5
months


+ASCVD

Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥ 45 years and with a BMI ≥ 27 kg/m²



Semaglutide 2.4 mg ⁴

Semaglutide treatment was associated with relatively lower risk of 5P-MACE^a



Event rate

Semaglutide group: 1.4%
Non-semaglutide group: 2.3%
HR: 0.64 (0.51–0.79)

Incidence of 5P-MACE per 1000 person-years

Semaglutide	12.8
Non-semaglutide	20.2
Absolute risk reduction	7.4

Individuals at risk (event rate [%])

Semaglutide group	7113 (–)	6557 (0.55)	5705 (0.75)	4856 (0.84)	4133 (1.05)	3365 (1.24)	2490 (1.35)	1169 (–)
Non-semaglutide group	21,339 (0.04)	19,654 (0.67)	17,052 (1.09)	14,410 (1.54)	12,246 (1.83)	9986 (2.04)	7416 (2.15)	3502 (–)

Shaded areas represent the 95% CIs. ^a5P-MACE was a composite of MI, stroke, hospitalization for HF, coronary revascularization and all-cause mortality. 5P-MACE, 5-point major adverse cardiovascular events; CI, confidence interval; HF, heart failure; HR, hazard ratio; MI, myocardial infarction.

Maximum follow-up was 23.5 months

Naltrexone/
Bupropion
(not recommended)¹⁵

+ASCVD

STEER Study Design

US Komodo Research Database

Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥ 45 years and with a BMI ≥ 27 kg/m²



Study population

- ≥ 45 years of age
- Overweight/obesity*
- Established ASCVD†
- Without diabetes
- Initiated semaglutide or tirzepatide on or after May 13, 2022

Jan 1, 2016*

Index date = treatment initiation

Jan 31, 2025

End of follow-up‡

Semaglutide 2.4 mg⁴

The propensity score model matched semaglutide and tirzepatide patients 1:1

Primary outcome measures:

- **Revised 3-point MACE:** Myocardial infarction, stroke, and **all-cause mortality**
- **Revised 5-point MACE:** Myocardial infarction, stroke, hospitalization for heart failure, coronary revascularization, and **all-cause mortality**

*Defined as body mass index ≥ 27 kg/m². †Defined as a diagnosis of myocardial infarction or ischemic stroke and/or evidence of peripheral artery disease. ‡End of follow-up for each patient was defined as the earliest of the end of the study period (Jan 31, 2025), end of continuous enrolment, initiation of a non-index GLP-1 or GLP-1/GIP receptor agonist, bariatric surgery, or death. ASCVD, atherosclerotic cardiovascular disease; GIP, glucose-dependent insulintropic polypeptide; GLP-1, glucagon-like peptide-1; MACE, major adverse cardiovascular event.

Wilson L, et al. Presented at the European Society of Cardiology (ESC) Congress together with World Congress of Cardiology; Madrid, Spain; August 29 - September 1, 2025.

Naltrexone/
Bupropion
(not recommended)¹⁵

+ASCVD

Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥ 45 years and with a BMI ≥ 27 kg/m²

Semaglutide 2.4 mg ⁴

Naltrexone/
Bupropion
(not recommended)¹⁵

Revised 3-Point MACE*


(Per-Protocol Analysis)

Per-protocol sensitivity analysis censored patients at treatment discontinuation (gap in therapy >30 days)

	Events, n (%)	Incidence rate per 1000 patient-years (95% CI)	Mean (SD) follow-up
Tirzepatide	39 (0.4)	10.3 (7.13-13.99)	4.3 (4.2)
Semaglutide	15 (0.1)	4.4 (2.34-7.03)	3.8 (4.1)

HR, 0.43 (95% CI, 0.24-0.78); $P = 0.005$

57%



*Revised 3-point MACE includes myocardial infarction, stroke, and all-cause mortality.
CI, confidence interval; HR, hazard ratio; MACE, major adverse cardiovascular event; SD, standard deviation.

Wilson L, et al. Presented at the European Society of Cardiology (ESC) Congress together with World Congress of Cardiology; Madrid, Spain; August 29 - September 1, 2025.

+ASCVD

Cardiovascular disease was defined as prior myocardial infarction, prior stroke, or clinically symptomatic peripheral arterial disease in individuals aged ≥45 years and with a BMI ≥27 kg/m²


Semaglutide 2.4 mg ⁴

Naltrexone/
Bupropion
(not recommended)¹⁵

Revised 5-Point MACE* (Per-Protocol Analysis)

Per-protocol sensitivity analysis censored patients at treatment discontinuation (gap in therapy >30 days)

	Events, n (%)	Incidence rate per 1000 patient-years (95% CI)	Mean (SD) follow-up
Tirzepatide	80 (0.8)	21.1 (16.63-26.14)	4.3 (4.2)
Semaglutide	42 (0.4)	12.3 (8.79-16.41)	3.8 (4.1)



*Revised 5-point MACE includes myocardial infarction, stroke, hospitalization for heart failure, coronary revascularization, and all-cause mortality. CI, confidence interval; HR, hazard ratio; MACE, major adverse cardiovascular event; SD, standard deviation.

Clinical heart failure (NYHA II–IV) with preserved or mildly reduced ejection fraction (LVEF $\geq 50\%$ for tirzepatide, $\geq 45\%$ for semaglutide), BMI ≥ 30 kg/m², impaired functional capacity (KCCQ-CSS < 80 –90), 6MWT distance ≥ 100 m, and elevated filling pressures (e.g., NT-proBNP, echocardiographic abnormalities, or recent HF hospitalization). Type 2 diabetes was an inclusion criterion in the tirzepatide trial.



Tirzepatide
15 mg¹¹


Semaglutide
2.4 mg⁵

Naltrexone/
Bupropion
(not recommended)¹⁵

MACE (A), HF composite (B), CV death (C), All-cause death (D) by HF subtype

- Semaglutide reduced MACE in both subtypes: **HR 0.65 (HFrEF)** and **0.69 (HFpEF)**; no interaction ($p = 0.97$).
- Semaglutide lowered the **HF composite endpoint** regardless of ejection-fraction category.
- Reductions in CV death and all-cause mortality were **directionally similar**, with a numerically greater absolute benefit in HFrEF due to higher baseline risk.




Cumulative incidence curves showing the risk of MACE comparing semaglutide with placebo according to presence or absence of HF. The cumulative incidence rate is calculated using the Aalen–Johansen method. CI, confidence interval; CV, cardiovascular; HF, heart failure; HR, hazard ratio; MACE, major adverse cardiovascular events.
Deanfield J, et al. *Lancet*. 2024;404(10454):773–786.





Hyperfiltration and Obesity-Related Glomerulopathy

↑ plasma flow → afferent vasodilation + efferent vasoconstriction → ↑ intraglomerular pressure
↑ sodium reabsorption → impaired tubulo-glomerular feedback
↑ RAAS & sympathetic activation (leptin), ↓ adiponectin, inflammation,



CKD

eGFR <60 mL/min/1.73 m² OR albuminuria (ACR ≥3.0 mg/mmol [≥30 mg/g])



Semaglutide 2.4 mg⁵


Tirzepatide 15 mg^{11,19}

Use if eGFR >30 mL/min/1.73 m²; exercise caution when eGFR <30 mL/min/1.73 m² due to limited data.

Orlistat (caution in obesity due to risks of nephrolithiasis or AKI)¹⁸

**Naltrexone/
Bupropion
(not recommended)¹⁵**

Time to first occurrence of the main 5-component kidney composite endpoint[†]



Semaglutide 2.4 mg reduced the risk of the main kidney composite endpoint by **22%** compared with placebo

Data are the observed (i.e. as measured) probability of patients experiencing their first occurrence of the main 5-component kidney composite endpoint during the in-trial period, analysed using the Kaplan–Meier method, and the estimated HR, analysed using a Cox regression model. Tied events were handled using the Exact method, if possible, or Efron’s method, if not. Numbers below the graph are the number of patients at risk. p values are two-sided and not adjusted for multiplicity

[†]The main 5-component kidney composite endpoint included death from kidney causes, initiation of chronic kidney replacement therapy (dialysis or transplantation), onset of persistent eGFR <15 mL/min/1.73 m², persistent ≥50% reduction in eGFR compared with baseline or onset of persistent macroalbuminuria. CI, confidence interval; eGFR, estimated glomerular filtration rate; HR, hazard ratio.

Colhoun HM et al. Nat Med 2024;30:2058–2066.

CKD

eGFR <60 mL/min/1.73 m² OR albuminuria (ACR ≥3.0 mg/mmol [≥30 mg/g])



Changes in eGFR over time

Semaglutide 2.4 mg⁵
Tirzepatide 15 mg^{11,19}

Use if eGFR >30 mL/min/1.73 m²; exercise caution when eGFR <30 mL/min/1.73 m² due to limited data.

Orlistat (caution in obesity due to risks of nephrolithiasis or AKI)¹⁸

Naltrexone/Bupropion (not recommended)¹⁵



Data are estimated mean (CI) changes from the estimated baseline value in eGFR, analysed using a mixed model for repeated measurements.

Numbers below the graphs are the number of patients contributing to the analysis.

p values are two-sided and not adjusted for multiplicity. Darker lines are used for the larger subgroups.

[†]Given gradual entry to the trial across the enrolment period and variable follow-up duration, data at 156 and 208 weeks are sparser compared with prior time points.

CI, confidence interval; eGFR, estimated glomerular filtration rate.

Colhoun HM et al. Nat Med 2024;30:2058–2066.

CKD

eGFR <60 mL/min/1.73 m² OR albuminuria (ACR ≥3.0 mg/mmol [≥30 mg/g])

Semaglutide 2.4 mg⁵
Tirzepatide 15 mg^{11,19}

Use if eGFR >30 mL/min/1.73 m²; exercise caution when eGFR <30 mL/min/1.73 m² due to limited data.


Orlistat (caution in obesity due to risks of nephrolithiasis or AKI)¹⁸

Naltrexone/Bupropion (not recommended)¹⁵

Changes in UACR over time

In the overall population

Treatment effect at 104 weeks: **-10.7%** (95% CI -13.2, -8.2; p<0.001)



Data are estimated mean (CI) changes from the estimated baseline value in UACR, analysed using a mixed model for repeated measurements. The change in UACR was analysed as the estimated mean ratio to baseline; for ease of interpretation, these ratios have been converted to relative percentage changes from baseline using the formula (estimated ratio - 1) × 100. Numbers below the graphs are the number of patients contributing to the analysis. p values are two-sided and not adjusted for multiplicity. Darker lines are used for the larger subgroups.
[†]Given gradual entry to the trial across the enrolment period and variable follow-up duration, data at 156 and 208 weeks are sparser compared with prior time points. CI, confidence interval; UACR, urinary albumin-to-creatinine ratio.
Colhoun HM et al. Nat Med 2024;30:2058–2066.


By subgroup

Treatment effect at 104 weeks (by UACR subgroups, mg/g)

<30: **-8.1%** (95% CI -10.6, -5.6; p<0.001)

30 to <300: **-27.2%** (95% CI -35.3, -18.1; p<0.001)

≥300: **-31.4%** (95% CI -54.9, 4.3; p=0.08)



UACR (mg/g):
Semaglutide 2.4 mg
Placebo



GLYCEMIC CONTROL

PREDIABETES

HbA1c 5.7–6.4%, fasting plasma glucose 100–125 mg/dL, or 2-hours OGTT glucose 140–199 mg/dL

- Tirzepatide 15 mg²
- Semaglutide 2.4 mg²²
- Liraglutide 3 mg^{12,74}
- Orlistat 120 mg¹⁴

Metformin: off-label use; may reduce the incidence of type 2 diabetes²⁴.

TYPE 2 DIABETES

HbA1c ≥ 6.5%, fasting plasma glucose ≥ 126 mg/dL, or 2-hours OGTT plasma glucose ≥ 200 mg/dL

- Tirzepatide 15 mg³¹
- Semaglutide 2.4 mg²⁹
- Liraglutide 3 mg³³

SGLT2 inhibitors support weight loss^{37,38,39,40,41}. If glycemic control allows, insulin^{44,45}, thiazolidinediones, and sulfonylureas should be avoided due to their propensity to induce weight gain.²⁰³



TYPE 1 DIABETES

(OFF-LABEL only for weight loss purposes)

- Tirzepatide 15 mg⁵²
- Semaglutide 2.4 mg⁵¹

Reversion to normoglycaemia

Primary endpoint (week 52) and exploratory endpoint (week 80)



Normoglycaemia was defined as having both HbA_{1c} <6.0% and FPG <5.5 mmol/L. Observed proportions of participants, during the in-trial observation period in the FAS. ORs are for the treatment policy estimand. CI, confidence interval; FAS, full analysis set; FPG, fasting plasma glucose; HbA_{1c}, glycated haemoglobin; OR, odds ratio. McGowan, B et al. *Obes Facts* 2024;17(suppl 1):72

Progression to T2D

Exploratory endpoints

GLYCEMIC CONTROL

PREDIABETES

HbA1c 5.7–6.4%, fasting plasma glucose 100–125 mg/dL, or 2-hours OGTT glucose 140–199 mg/dL

- Tirzepatide 15 mg ²
- Semaglutide 2.4 mg ²²
- Liraglutide 3 mg ^{12,74}
- Orlistat 120 mg ¹⁴

Metformin: off-label use; may reduce the incidence of type 2 diabetes²⁴.

TYPE 2 DIABETES

HbA1c ≥ 6.5%, fasting plasma glucose ≥ 126 mg/dL, or 2-hours OGTT plasma glucose ≥ 200 mg/dL

- Tirzepatide 15 mg ³¹
- Semaglutide 2.4 mg ²⁹
- Liraglutide 3 mg ³³


SGLT2 inhibitors support weight loss ^{37,38,39,40,41}. If glycemic control allows, insulin^{44,45}, thiazolidinediones, and sulfonylureas should be avoided due to their propensity to induce weight gain.²⁰³

TYPE 1 DIABETES


(OFF-LABEL only for weight loss purposes)

- Tirzepatide 15 mg ⁵²
- Semaglutide 2.4 mg ⁵¹


Progression to T2D at week 52





Progression to T2D at week 80
Off treatment



T2D was defined as having HbA_{1c} ≥6.5% and/or FPG ≥7.0 mmol/L verified with a repeated blood sample within 4 weeks. Observed proportions of participants, during the in-trial observation period in the FAS. FAS, full analysis set; FPG, fasting plasma glucose; HbA_{1c}, glycated haemoglobin; T2D, type 2 diabetes. McGowan et al. presented at the 2024 31st European Congress on Obesity, Venice, Italy, 12–15 May 2024. Abstract AS12.02



Liver fibrosis stage is a predictor of increased mortality



Meta-analysis of 5 studies reporting fibrosis stage-specific mortality, N=1495.
PYF, patient-years of follow-up.
Dulai PS et al. *Hepatology*. 2017;65:1557–65.

MASLD/ MASH

Adults with biopsy-confirmed MASH, fibrosis stage F2–F3 for semaglutide 2.4 mg and tirzepatide 15 mg; F0–F3 for liraglutide 3 mg. BMI ≥ 25 kg/m² for semaglutide 2.4 mg and liraglutide 3 mg; BMI ≥ 27 kg/m² for tirzepatide 15 mg, with or without type 2 diabetes; BMI not restricted, and with or without type 2 diabetes, for semaglutide 2.4.





Semaglutide 2.4 mg⁵⁴

Tirzepatide 15 mg⁵⁶

Liraglutide 3 mg OD⁵⁷

Steatohepatitis resolution and improvement in liver fibrosis

Proportion of patients at Week 72 (full analysis set)



Significantly more patients with MASH F2–F3 treated with semaglutide 2.4 mg achieved **both primary endpoints of MASH resolution (62.9%) and improvement in liver fibrosis (37.0%)** than those treated with placebo (**34.1%, 22.5% respectively**)

Analysis set: FAS (interim), first 800 randomised subjects. EDP: Estimated difference in responder proportions with 95% confidence interval and two-sided p-value. *Resolution of steatohepatitis is defined as a NAS of 0–1 for inflammation, 0 for ballooning and any value for steatosis according to NASH CRN. No worsening of liver fibrosis is defined as no increase in fibrosis score. Fibrosis is graded on the NASH CRN fibrosis scale from 0–4. †Improvement in fibrosis is defined as ≥ 1 grade improvement on the NASH CRN fibrosis scale. No worsening of steatohepatitis is defined as no increase from baseline in NAS score for ballooning, inflammation or steatosis. The absolute difference between responder proportions, 95% confidence interval, P value was generated with the use of Cochran–Mantel–Haenszel (CMH) test stratified by baseline diabetes status (medical history) and fibrosis stage (eligibility read). #Missing data were handled by reference-based multiple imputation and Rubin's rule based on the Mantel–Haenszel estimator and Sato's estimate of the standard error (reference) were used to aggregate results. CI, confidence interval; EDP, estimated difference in responder proportions; F, fibrosis stage; MASH, metabolic dysfunction associated steatohepatitis; NASH CRN, Non-Alcoholic Steatohepatitis Clinical Research Network. Newsome PN et al. Oral Presentation at American Association for the Study of the Liver The Liver Meeting; Late Breaker 5018; November 19 2024; San Diego, USA.

Sanyal AJ, Newsome PN, Kliers I, et al. Phase 3 Trial of Semaglutide in Metabolic Dysfunction-Associated Steatohepatitis. *N Engl J Med*. 2025;392(21):2089-2099


MASLD/ MASH

FDA Approves Treatment for Serious Liver Disease Known as 'MASH'

Action Will Provide New Therapy for Growing Public Health Issue

Adults with biopsy-confirmed MASH, fibrosis stage F2–F3 for semaglutide 2.4 mg and tirzepatide 15 mg; F0–F3 for liraglutide 3 mg. BMI ≥ 25 kg/m² for semaglutide 2.4 mg and liraglutide 3 mg; BMI ≥ 27 kg/m² for tirzepatide 15 mg, with or without type 2 diabetes; BMI not restricted, and ,with or without type 2 diabetes, for semaglutide 2.4.

Resolution of steatohepatitis and improvement in liver fibrosis



Significantly more patients with MASH F2–F3 treated with semaglutide 2.4 mg achieved resolution of steatohepatitis and improvement in liver fibrosis (32.8%) than those treated with placebo (16.2%)

Analysis set: FAS (interim), first 800 randomised subjects. Metabolic dysfunction-associated steatohepatitis (MASH) was previously known as nonalcoholic steatohepatitis (NASH). EDP: Estimated difference in responder proportions with 95% confidence interval and two-sided p-value. Resolution of steatohepatitis is defined as a NAS of 0-1 for inflammation, 0 for ballooning and any value for steatosis according to NASH CRN. Improvement in fibrosis is defined as ≥ 1 grade improvement on the NASH CRN fibrosis scale. NASH CRN: Non-Alcoholic Steatohepatitis Clinical Research Network. The absolute difference between responder proportions, 95% confidence interval, P value was generated with the use of Cochran -Mantel-Haenszel (CMH) test stratified by baseline diabetes status (medical history) and fibrosis stage (eligibility read). Missing data were handled by reference-based multiple imputation and Rubin's rule based on the Mantel-Haenszel estimator (reference) and Sato's estimate of the standard error were used to aggregate results. CI, confidence interval; EDP, estimated difference in responder proportions; F, fibrosis stage; MASH, metabolic dysfunction-associated steatohepatitis; n, number of participants. Newsome PN et al. Oral Presentation at American Association for the Study of the Liver The Liver Meeting; Late Breaker 5018; November 19 2024; San Diego, USA.

Sanyal AJ, Newsome PN, Kliers I, et al. Phase 3 Trial of Semaglutide in Metabolic Dysfunction-Associated Steatohepatitis. *N Engl J Med*. 2025;392(21):2089-2099

Semaglutide 2.4 mg⁵⁴
Tirzepatide 15 mg⁵⁶
Liraglutide 3 mg OD⁵⁷

OBESEITY TREATMENT BASED ON PATIENT PHENOTYPING

FIRST-LINE THERAPY IS LIFESTYLE MODIFICATION (MEDICAL NUTRITIONAL APPROACH AND IMPLEMENTATION OF PHYSICAL ACTIVITY AND BEHAVIOURAL THERAPY)*

EXCLUDE ENDOCRINE FORMS OF OBESITY, if suspected, investigate MONOGENIC FORMS

ASSESSMENT OF THE PRESENCE OF COMPLICATIONS AND PATIENT PHENOTYPING

OBESEITY TREATMENT BASED ON PATIENT COMPLICATIONS AND PHENOTYPING

METABOLIC CARDIO RENAL COMPLICATIONS						MECHANICAL COMPLICATIONS		WEIGHT LOSS TARGETS /OWCMC**	AGE	EATING BEHAVIOUR	SPECIAL POPULATIONS	COSTS /QALY
+ASCVD	Indicators of High CV Risk:	HF	CKD	GLYCEMIC CONTROL	MASH/MASLD	OSTEOARTHRITIS	OSAS					
Cardiovascular disease was defined as a previous myocardial infarction, previous stroke, or symptomatic peripheral arterial disease in people who are ≥ 45 years of age, with a BMI ≥ 27 kg/m².	Definitions vary; most comprise ≥ 40 years of age, BMI ≥ 27 kg/m², male gender, high blood pressure, atherogenic dyslipidemia, smoking, prediabetes, ASCVD risks (ACC/AHA).	Clinical heart failure (NYHA II–IV) with LVEF ≥50% (tirzepatide) or ≥45% (semaglutide), BMI ≥30 kg/m², reduced functional capacity (KCCQ-CSS <80 or <90), 6-minute walk ≥100 m, and elevated filling pressures (NT-proBNP, echocardiographic criteria, or recent HF hospitalization). T2D was all in the tirzepatide trial.	eGFR < 60 ml/ min per 1.73 m² OR albuminuria (ACR ≥ 3.0 mg/mmol (30mg/g)	PREDIABETES (HbA1c ≥ 5.7% and < 6.5%, FPG > 100 ng/dl < 126 mg/dl, OGTT 120 min glucose ≥ 140 mg/dl < 200 mg/dl) Tirzepatide 15 mg ² Semaglutide 2.4 mg ²² Liraglutide 3 mg ^{12,74} Orlistat 120 mg ¹⁴ Metformin: off-label; may reduce diabetes incidence ²⁴ .	Adults with biopsy-confirmed MASH, fibrosis stage (F2–F3) for semaglutide 15 mg, F0–F3 for liraglutide 3 mg, BMI ≥25 kg/m² for semaglutide 2.4 mg and liraglutide 3 mg, BMI ≥27 kg/m² for tirzepatide 15 mg; with or without type 2 diabetes but not for semaglutide 2.4.	Adults ≥18 years with BMI ≥30 kg/m², clinically and radiographically confirmed knee osteoarthritis (ACR criteria; Kellgren–Lawrence grade 2–3), and baseline knee pain ≥40/100 on the WOMAC pain scale.	Adults ≥18 years, BMI ≥30 kg/m², with moderate-to-severe obstructive sleep apnea (AHI ≥15 events/hour); CPAP use allowed (tirzepatide) or excluded (liraglutide); no diagnosis of T2D.	Set individualized weight management goals 5-10% WL Orlistat ¹⁴ Liraglutide 3 mg ⁷⁴ Naltrexone/ Bupropion ²⁷ Phentermine/Tipiramate (low-mid doses) ⁶⁰	≥ 2 y.o. ¹⁰⁴ Setmelanotide (rare genetic obesity) Metreleptin (rare genetic obesity) 6-12 y.o. Liraglutide 3mg ¹⁰⁹ (EMA only) 12-18 y.o. Liraglutide 3mg ¹¹² Semaglutide 2.4mg ¹¹³ Phentermine/Topiramate ¹¹³ (FDA only) Orlistat 120 mg ¹¹⁴ (FDA only) 18-65 y.o. Phentermine/Topiramate ¹²⁶ (FDA decision): Tirzepatide 15 mg Semaglutide 2.4 mg Naltrexone/Bupropion ^{27,28} Phentermine/Topiramate ²⁶ Orlistat ¹⁴ >65 y.o. Semaglutide 2.4 mg ³ (Usable; no dose adjustment necessary) Tirzepatide 15 mg ³ (Usable; monitor sensitivity in elderly) Liraglutide 3 mg ¹²⁴ (Usable; limited data >75 yrs) Naltrexone/Bupropion ²⁷ (Caution; limited data ≥65 yrs) Phentermine/Topiramate ²⁶ (Caution; monitor CNS/renal) Orlistat ¹⁴ (Usable; limited efficacy data) Sarcopenic obesity Supervised Resistance Training + Protein Intake ≥1.2–1.5 g/kg/day ¹¹⁸ Liraglutide 3 mg is preferred for gradual weight loss with lean mass preservation and proven cardiovascular safety ³³	EMOTIONAL EATING Bupropion / Naltrexone ¹⁴⁸ Tirzepatide 15 mg ¹²⁵ Semaglutide 2.4 mg ¹²⁵ BINGE EATING Not AOMs approved explicitly for binge eating disorder. Liraglutide 3.0 mg ¹²³ Phentermine/Topiramate ¹²⁸ Naltrexone/Bupropion ²⁸ Orlistat 120 mg ¹²⁹ Semaglutide 2.4 mg ¹²⁵ Tirzepatide 15 mg ^{128,134}	CANCER safety No significant obesity-related cancer risk from obesity-related RCT Semaglutide 2.4 mg ¹⁴⁷ Tirzepatide 15 mg ¹⁴⁷ Evidence from meta-analysis suggests long-term GLP-1 RA use may increase thyroid cancer risk in mixed T2D populations. ¹⁴⁵ No evidence of increased cancer risk (for other AOMs: Orlistat, Phentermine/Topiramate, Bupropion/Naltrexone) Familial Partial Lipodystrophy 1 -3 (FPLD1-3) Acquired Partial Lipodystrophy (Barraquer-Simons) HLM Associated Lipodystrophy Semaglutide 2.4 mg ¹⁵² MONOGENIC OR SYNDROMIC OBESITY Congenital leptin deficiency Metreleptin ¹⁵³ (biallelic POMC, PCSK1, or LEPR deficiency and for BBS) Setmelanotide 3 mg ¹⁵⁴ ACQUIRED HYPOTALAMIC OBESITY Setmelanotide 3 mg ¹⁵⁴ Semaglutide 2.4 mg ¹⁵⁵ PREGNANCY/WILLINGNESS TO CONCEIVE AOMs are either contraindicated or advised against during pregnancy due to insufficient safety data and potential risks to the fetus Stop Liraglutide 3 mg/Semaglutide 2.4 mg at least 2 months before Stop Tirzepatide 15 mg at least 1 month before	BS 176,177
Semaglutide 2.4 mg	Tirzepatide 15 mg ^{2,10} Semaglutide 2.4 mg ^{3,4} Liraglutide 3 mg ¹²	Tirzepatide 15 mg ¹¹ Semaglutide 2.4 mg ⁵	Semaglutide 2.4 mg ⁵ Tirzepatide 15 mg ^{11,19} (use if eGFR>30 ml/min; If eGFR <30 use with caution due to limited data) Orlistat (caution in obesity due to risks of nephrolithiasis or AKI) ¹⁸	Tirzepatide 15 mg ¹ Semaglutide 2.4 mg ²⁹ Liraglutide 3 mg ³³ SGLT2is support weight loss ^{37,38,39,40,41} ; if glycemic control allows, insulin ^{44,45} , thiazolidinediones, and sulfonylureas should be avoided ²⁰³ TYPE 1 DIABETES (OFF-LABEL only for weight loss purposes) Tirzepatide 15 mg ⁵² Semaglutide 2.4 mg ⁵¹	Semaglutide 2.4 mg ⁴ Tirzepatide 15 mg ⁵ Liraglutide 3 mg OD ⁵⁷	Semaglutide 2.4 mg ⁶³ Tirzepatide 15 mg ⁶³ Orlistat 120 mg ¹⁴ Naltrexone/bupropione ⁶⁴ Phentermine/Topiramate ²⁶	Tirzepatide 11 mg ⁶⁸ Liraglutide 3 mg ⁶⁹ Phentermine/Topiramate ⁷⁰	10-20% WL Tirzepatide 5 mg ² Semaglutide 2.4mg ⁴ Phentermine/Topiramate ²⁶ (low-mid doses) ⁶⁰ >20% WL Tirzepatide 15mg ² Semaglutide 7.2 mg ⁷⁷ (not available yet) Cagrisema ⁷⁸ (not available yet)				
Naltrexone/ Bupropion (not recommended) ¹⁵	Naltrexone/ Bupropion (not recommended) ¹⁵	Naltrexone/ Bupropion (not recommended) ¹⁵	Naltrexone/ Bupropion (not recommended) ¹⁵									

If additional weight loss and reduction of weight-related complications are needed

Bariatric Surgery¹⁷⁷

If insufficient weight loss or weight regain after BS consider: Tirzepatide¹⁹⁷, Semaglutide¹⁹⁴, Liraglutide^{195,196}

*This framework promotes a phenotype-based, individualized approach to obesity care, prioritizing complications and patient preferences over rigid treatment hierarchies, with patient-HCP shared decision-making considering goals, tolerability, contraindications, and cost. **OWCMC, obesity without clinically manifest complications